

LAYERED XY-MODELS, ANYON SUPERCONDUCTORS, AND SPIN-LIQUIDS

J. P. Rodriguez

*Dept. of Physics and Astronomy, California State University, Los Angeles, CA 90032**
and Theoretical Division, Los Alamos National Laboratory, Los Alamos, NM 87545.

Abstract

The partition function of the double-layer XY model in the (dual) Villain form is computed exactly in the limit of weak coupling between layers. Both layers are found to be locked together through the Berezinskii-Kosterlitz-Thouless transition, while they become decoupled well inside the normal phase. These results are recovered in the general case of a finite number of such layers. When re-interpreted in terms of the dual problems of lattice anyon superconductivity and of spin-liquids, they also indicate that the essential nature of the transition into the normal state found in two dimensions persists in the case of a finite number of weakly coupled layers.

PACS Indices: 75.10.Hk, 74.20.Kk, 74.20.Mn, 11.15.Ha

* Permanent address.

The layered crystal structure common to high-temperature superconductors is known to be reflected in their electronic properties.¹ There are suggestions in the theoretical literature that the effect of strong repulsive interactions present within a single such copper-oxygen layer result in exotic anyonic superconducting states and/or spin-liquid states.^{2–5} Recently, the author has shown that both of these groundstates undergo a phase transition into a normal phase at non-zero temperature, where the corresponding particle-hole excitations become deconfined upon heating through the transition.⁶ The transition itself is dual to that of the two-dimensional (2D) Coulomb gas.^{7,8} In the present context of oxide superconductivity, therefore, one is naturally led to the question of how a finite number of such weakly coupled layers behave.

In response to this query, we shall study the problem of a finite number of weakly coupled layers of square-lattice XY -models in the Villain form,⁹ which is known to lead to a dual theory of (layered) compact quantum electrodynamics (QED) in the strong-coupling limit.¹⁰ The Abelian gauge-field that appears here describes chiral spin-fluctuations in the case of the 2D spin-liquid state, while it represents the statistical flux-tube attached to each psuedo-fermion in the case of anyon superconductivity.^{3–5,11} Korshunov has given an approximate solution of the XY -model with infinitely many weakly coupled layers, where he finds a Berezinskii-Kosterlitz-Thouless (BKT) transition within each layer, followed by an inter-layer decoupling transition at much higher temperature that is characterized by the binding of vortex rings (fluxons) lying in between consecutive layers.¹² Note, therefore, that the lower-temperature superfluid BKT transition involves the *entire* three-dimensional (3D) lattice in this scenario, which is consistent with previous Monte Carlo simulations.¹³ In this Letter, we first solve the weakly coupled double-layer XY model in the Villain form *exactly*, recovering Korshunov’s approximate results for the case of infinitely many layers in the processes. The intra-layer phase auto-correlation function of this double-layer XY -model is computed, where we find that it falls off algebraically with distance below the BKT transition, as expected. Notably, the exponent of the correlation function in question vanishes linearly with temperature, yet the coefficient is only half as big as that of the single-layer case.⁹ This indicates that the existence of neighboring layers enhances long-range phase correlations, as expected. In addition, the inter-layer phase autocorrelation function is computed, where we find that it coincides with the prior intra-layer phase

autocorrelation function at temperatures below the decoupling transition. This, on the other hand, gives strong support to the claim that the double-layers become locked together at temperatures below the latter transition. Last, we show that the above double-layer results persist in the general case of a finite number of weakly coupled layers, suggesting that the superfluid transition exhibited by the XY model with a finite number of layers is 3D-like.^{12,13}

When the above results are re-interpreted in terms of the dual problem of double-layer anyon superconductivity and spin-liquids, they yield that a confining string exists between particle-hole excitations at temperatures below the deconfinement transition.¹⁰ Note that the consecutive layers have already become decoupled at this stage via the inter-layer fluxon-antifluxon unbinding transition that occurs at much lower temperature in this case. Taking into account arguments given previously by the author,⁶ this implies that the basic character of the single-layer particle-hole deconfinement transition into the normal state persists in the case of double-layer anyon superconductors and spin-liquids; i.e., the transition remains 2D.

Let us first consider the generalized correlation function $\langle \exp[i \sum_r p(r) \phi(r)] \rangle = Z[p]/Z[0]$ of the N -layered XY -model, where the partition function $Z[p] = \int \mathcal{D}\phi(r) e^{-E/T}$ has an energy functional

$$\begin{aligned} \frac{E}{T} = & \beta_{\parallel} \sum_{l=1}^N \sum_{\vec{r}} \sum_{\mu=x,y} \{1 - \cos[\Delta_{\mu} \phi(\vec{r}, l)]\} + \beta_{\perp} \sum_{l=1}^{N-1} \sum_{\vec{r}} \{1 - \cos[\phi(\vec{r}, l) - \phi(\vec{r}, l+1)]\} \\ & - i \sum_{l=1}^N \sum_{\vec{r}} p(\vec{r}, l) \phi(\vec{r}, l). \end{aligned} \quad (1)$$

Here $\vec{r}$ ranges over the square-lattice and l denotes the layer index, while $\beta_{\parallel, \perp} = J_{\parallel, \perp}/k_B T$ is proportional to nearest-neighbor coupling-constants satisfying $J_{\parallel} \gg J_{\perp}$. Also, $\Delta_{\mu} \phi(r) = \phi(r + \hat{\mu}) - \phi(r)$ is the lattice difference operator. The particular correlation to be probed is determined by the fixed integer field $p(\vec{r}, l)$; e.g., the choice $p(\pm r/2, 0, 1) = \pm 1$ corresponds to the intra-plane auto-correlation function. After making the usual (low-temperature) Villain substitution for the exponential above,⁹ $e^{-\beta[1 - \cos \theta]} \rightarrow (2\pi\beta)^{-1/2} \sum_{n=-\infty}^{\infty} e^{in\theta} e^{-n^2/2\beta}$, and then integrating over the phase-field, we obtain the following dual representation equiv-

alent to N -layered compact QED in the strong-coupling limit (modulo a constant):¹⁰

$$Z[p] = \sum_{\{n_\mu(r)\}} \Pi_r \delta \left[\Delta_\nu n_\nu|_r - \sum_{r'} p(r') \delta_{r'r} \right] \exp \left[-\frac{1}{2\beta_\parallel} \sum_{l=1}^N \sum_{\vec{r}} \vec{n}^2(\vec{r}, l) - \frac{1}{2\beta_\perp} \sum_{l=1}^{N-1} \sum_{\vec{r}} n_z^2(\vec{r}, l) \right], \quad (2)$$

where $n_\mu(r)$ is an integer link-field on the layered lattice structure of points $r = (\vec{r}, l)$, with $\mu = x, y, z$ and $\vec{n} = (n_x, n_y)$. In the context of the dual systems of layered anyon superconductors and spin-liquids, $\vec{n}(\vec{r}, l)$ represents the confining statistical electric field experienced between particle-hole excitations about such groundstates.⁶ Here one must make the identification $\beta_{\perp, \parallel}^{-1} \rightarrow g_{\perp, \parallel}^2 \hbar \omega_0 / k_B T'$, where $g_{\perp, \parallel}$ denote the corresponding gauge-field coupling constants, and where T' is the temperature in these dual systems. The Debye frequency scale, $\omega_0 = c_0 a^{-1}$, is set by the characteristic velocity c_0 , which represents the spin-wave velocity in the case of spin-liquids, and the zero-sound speed in the case of anyon superconductivity.⁶ Note that the case of weakly coupled planes, $J_\perp \ll J_\parallel$, also corresponds to weakly electro-magnetically coupled layers in the dual system, where the coupling-constants satisfy $g_\perp^{-2} \ll g_\parallel^{-2}$.

Double-layer. Consider now the dual representation outlined above for the special case of two coupled XY-model/compact QED layers. This problem may be solved exactly in the limit of weak interlayer coupling, $J_\perp \ll J_\parallel$, by first making the decomposition $\vec{n}(\vec{r}, 1) = \vec{n}'(\vec{r}, 1) - \vec{n}_-(\vec{r})$ and $\vec{n}(\vec{r}, 2) = \vec{n}'(\vec{r}, 2) + \vec{n}_-(\vec{r})$, such that the intra-layer fields $\vec{n}'(\vec{r}, l)$ satisfy $\vec{\nabla} \cdot \vec{n}'|_r = \sum_{r'} p(r') \delta_{rr'}$, while the inter-layer field $\vec{n}_-(\vec{r})$ satisfies $\vec{\nabla} \cdot \vec{n}_-|_{\vec{r}} = n_z(\vec{r})$, with $\vec{\nabla} = (\Delta_x, \Delta_y)$. We now take the customary potential representation $\vec{n}_- = -\vec{\nabla} \Phi$ for the inter-layer field, which yields $\Phi(\vec{r}) = \sum_{\vec{r}'} G^{(2)}(\vec{r} - \vec{r}') n_z(\vec{r}')$, where $G^{(2)}(\vec{r}) = (2\pi)^{-2} \int_{\text{BZ}} d^2 k (e^{i\vec{k} \cdot \vec{r}} - 1) [4 - 2\cos(k_x a) - 2\cos(k_y a)]^{-1}$ is the Greens function for the square lattice. Note that fluxon-charge neutrality is presumed;⁹ i.e., $\sum_{\vec{r}} n_z(\vec{r}) = 0$. The requirement that the original field $\vec{n}$ be integer valued indicates that the intra-layer field $\vec{n}'$ is not integer-valued, in general. However, in the presently considered limit of weak coupling, $J_\perp / J_\parallel \rightarrow 0$, the concentration of inter-layer (fluxon) charge, n_z , is exponentially small. This implies that the inter-layer field, $\vec{n}_-$, is equally small in magnitude, and that the intra-layer field, $\vec{n}'$, is very close to being integer valued. After making a suitable (lattice) integration by parts of the energy functional in Eq. (2), we thus obtain the factorization $Z = Z_{\text{DG}}^{(1)} Z_{\text{DG}}^{(2)} Z_{\text{CG}}$ for the partition function in the limit of weak coupling between layers, where the intra-layer factors represent 2D discrete gaussian models that

read

$$Z_{\text{DG}}^{(l)}[p] = \sum_{\{\vec{n}'(\vec{r}, l)\}} \Pi_{\vec{r}} \delta \left[\vec{\nabla} \cdot \vec{n}'|_{\vec{r}, l} - \sum_{\vec{r}'} p(\vec{r}', l) \delta_{\vec{r}', \vec{r}} \right] \exp \left[-\frac{1}{2\beta_{\parallel}} \sum_{\vec{r}} \vec{n}'^2(\vec{r}, l) \right], \quad (3)$$

and where the inter-layer factor is given by the 2D Coulomb gas ensemble

$$\begin{aligned} Z_{\text{CG}}[p] &= \sum_{\{n_z(\vec{r})\}} \exp \left\{ -\frac{1}{2\beta_{\parallel}} \sum_{\vec{r}} 2[n_z(\vec{r}) - p(\vec{r}, 1) + p(\vec{r}, 2)] \Phi(\vec{r}) - \frac{1}{2\beta_{\perp}} \sum_{\vec{r}} n_z^2(\vec{r}) \right\} \\ &= \sum_{\{n_z(\vec{r})\}} \exp \left\{ -\frac{1}{2\beta_{\parallel}} \sum_{\vec{r}, \vec{r}'} 2[n_z(\vec{r}) - p(\vec{r}, 1) + p(\vec{r}, 2)] G^{(2)}(\vec{r} - \vec{r}') n_z(\vec{r}') - \frac{1}{2\beta_{\perp}} \sum_{\vec{r}} n_z^2(\vec{r}) \right\}. \end{aligned} \quad (4)$$

Considering first the case of absent probe charges, $p(r) = 0$, we immediately see that each layer (3) undergoes a pure 2D XY /discrete-gaussian model transition at $k_B T_c \lesssim \frac{\pi}{2} J_{\parallel}$, while the inter-layer links $n_z(\vec{r})$ undergo an inverted 2D Coulomb gas binding transition at $k_B T_* = 4\pi J_{\parallel}$ in the limit of weak inter-layer coupling, $J_{\perp} \ll J_{\parallel}$. We therefore find that Korshunov's approximate results for the corresponding problem of infinitely many layers are *exact* in the present instance of double layers.¹² Following this author, the latter high-temperature transition therefore corresponds to an inter-layer decoupling transition mediated by the binding of vortex rings lying in between the double-layers.

Next consider the intra-layer phase autocorrelation function, $C_{11}(R) = \langle \exp[i\phi(0, 1) - i\phi(\vec{R}, 1)] \rangle$, for the XY -model, which corresponds to fixing two opposing unit charges separated by a distance R within the dual compact QED representation; e.g., we have $p(\vec{r}, 1) = \delta_{\vec{r}, 0} - \delta_{\vec{r}, \vec{R}}$ and $p(\vec{r}, 2) = 0$. By the previous factorization for the partition function, we may therefore write $C_{11}(R) = C_{\parallel}(R)C_{\perp}(R)$, where $C_{\parallel}(R) = Z_{\text{DG}}^{(1)}[p]/Z_{\text{DG}}^{(1)}[0]$ and $C_{\perp}(R) = Z_{\text{CG}}[p]/Z_{\text{CG}}[0]$. The “parallel” autocorrelation is given by that of the pure square-lattice XY model;⁹ i.e., $C_{\parallel}(R) = (r_0/R)^{\eta_{\parallel}} e^{-R/\xi_{\parallel}}$, where the correlation length diverges as $\xi_{\parallel}/a \sim \exp[A/(T/T_c - 1)^{1/2}]$ just above T_c ,⁸ and where the exponent is given by $\eta_{\parallel} = (2\pi\beta_{\parallel})^{-1}$ at low-temperature and by $\eta_{\parallel} = 1/4$ at T_c . Here, $\ln r_0^{-1} = 1.6169$. To compute the “perpendicular” factor $C_{\perp}(R)$ to the intra-layer correlation function, we first re-express the right side of Eq. (4) as $Z_{\text{CG}}[p]/Z_{\text{CG}}[0] = \exp\{[G_{\text{MF}}^{(2)}(R) - G^{(2)}(R)]/2\beta_{\parallel}\}$, where the meanfield renormalized Coulomb interaction is defined by⁷

$$G_{\text{MF}}^{(2)}(R) = 2\beta_{\parallel} \ln \left[(Z_{\text{CG}}[0])^{-1} \sum_{\{n_z(\vec{r})\}} \exp \left\{ -\frac{1}{2\beta_{\parallel}} \sum_{\vec{r}, \vec{r}'} 2 \left[n_z(\vec{r}) - \frac{1}{2} (\delta_{\vec{r}, 0} - \delta_{\vec{r}, \vec{R}}) \right] G^{(2)}(\vec{r} - \vec{r}') \times \right. \right.$$

$$\times \left[n_z(\vec{r}') - \frac{1}{2}(\delta_{\vec{r}',0} - \delta_{\vec{r}',\vec{R}}) \right] - \frac{1}{2\beta_\perp} \sum_{\vec{r}} n_z^2(\vec{r}) \Bigg\}. \quad (5)$$

Therefore, for $T < T_*$ the above meanfield renormalized Coulomb interaction is short-ranged, yielding $C_\perp(R) = \exp[-G^{(2)}(R)/2\beta_\parallel] \cong (R/r_0)^{1/4\pi\beta_\parallel}$, while it is dielectrically renormalized to $G_{\text{MF}}^{(2)}(R) = \epsilon^{-1}(T)G^{(2)}(R)$ at $T > T_*$, yielding $C_\perp(R) \cong (R/r_0)^{(1-\epsilon^{-1})/4\pi\beta_\parallel}$. Putting together all of these results, we obtain $C_{11}(R) = (r_0/R)^\eta e^{-R/\xi_\parallel}$ for the inter-layer correlator, where the temperature dependent exponent is given by

$$\eta = \begin{cases} \frac{1}{4\pi\beta_\parallel}; & T \ll T_c \\ \frac{1}{8}; & T \sim T_c \\ \eta_\parallel - \frac{1-\epsilon^{-1}}{4\pi\beta_\parallel}; & T > T_*. \end{cases} \quad (6)$$

Notice that the above correlations are strengthened by the second layer with respect to those of a single layer below the decoupling transition T_* , while they approach those of an isolated single layer at high-temperature [$\lim_{T \rightarrow \infty} \epsilon(T) = 1$].

To compute the inter-layer phase auto-correlation function $C_{12}(R) = \langle \exp[i\phi(0,1) - i\phi(\vec{R},2)] \rangle$, it becomes useful to consider the corresponding probe charges as a superposition of the previous intra-layer ones along with the minimum-distance inter-layer configuration $p'(\vec{r},1) = \delta_{\vec{r},\vec{R}}$ and $p'(\vec{r},2) = -\delta_{\vec{r},\vec{R}}$. The latter configuration can be incorporated into the inter-layer field $\vec{n}_-(\vec{r})$, resulting in the modified constraint $\vec{\nabla} \cdot \vec{n}_-|_{\vec{r}} = n_z(\vec{r}) - \delta_{\vec{r},\vec{R}}$. After appropriately modifying the manipulations beginning with Eq. (2), we ultimately obtain $C_{12}(R) = C_\parallel(R) \exp\{-[G^{(2)}(R) + G_{\text{MF}}^{(2)}(R)]/2\beta_\parallel\} = (r_0/R)^{\eta_\parallel - (1+\epsilon^{-1})/4\pi\beta_\parallel} e^{-R/\xi_\parallel}$ for the inter-layer correlation function. Notice, therefore, that this inter-layer correlator is *equal* to the intra-layer correlator, $C_{11}(R)$, at temperatures below the decoupling transition, $T < T_*$, where $\epsilon^{-1}(T) = 0$. We see explicitly, therefore, that the double-layers become locked at temperatures below T_* .

In terms of double-layer compact QED, which describes the gauge-field dynamics in both double-layer anyon superconductors and spin-liquids,⁶ the above discussion implies that there exists a deconfinement transition at $k_B T'_c \gtrsim \frac{2}{\pi} g_\parallel^2 \hbar \omega_0$, below which both intra-layer [$C_{11}(R)$] and inter-layer [$C_{12}(R)$] particle-hole excitation in either system are confined

by a statistical electric flux tube¹⁰ with string tension $\gamma(T') = k_B T' / \xi_{\parallel}(T')$. Note that this result is identical to that of a single layer. The present double-layer case, however, also shows a decoupling transition at $k_B T'_* = (4\pi)^{-1} g_{\parallel}^2 \hbar \omega_0 \sim k_B T'_c / 10$ between the corresponding in-plane statistical electric fields, $\vec{n}(\vec{r}, l)$. In particular, the Fourier transform of the inter-layer correlation function for this field is given by

$$\langle \vec{n}(\vec{k}, 1) \cdot \vec{n}(-\vec{k}, 2) \rangle = -\langle \vec{n}_-(\vec{k}) \cdot \vec{n}_-(-\vec{k}) \rangle = -\frac{k^2}{\epsilon_0(k^2 + \xi_{\perp}^{-1})}, \quad (7)$$

where $\xi_{\perp}(T')$ denotes the Debye screening length for the 2D Coulomb gas of fluxons. Above, we have used the potential representation $\vec{n}_- = -\vec{\nabla}\Phi$ for the “difference” field, along with the identity $\langle \Phi(\vec{k})\Phi(-\vec{k}) \rangle = \epsilon_0^{-1}(k^2 + \xi_{\perp}^{-1})^{-1}$ for the Coulomb gas correlation function. Hence, the homogeneous inter-layer correlation $\langle \vec{n}(0, 1) \cdot \vec{n}(0, 2) \rangle$ jumps from $-\epsilon_0^{-1}(T'_*) = -\frac{2}{\pi}$ to zero upon heating through decoupling transition-temperature T'_* . This indicates that the deconfinement transition at $T'_c > T'_*$ is 2D, since the layers have already become decoupled by then. Finally, we note that the specific-heat of the present double-layer model for strong-coupling compact QED naturally decouples into $C_v = 2C_{\text{DG}} + C_{\text{CG}}$, where C_{DG} and C_{CG} denote the contributions due to an isolated layer (3) described by the discrete-gaussian model⁶ and those due to the inter-layer Coulomb-gas ensemble (4), respectively. Given that $C_{\text{CG}}(T')$ shows a smooth peak just above⁸ T'_* , while $C_{\text{DG}}(T')$ exhibits one just below⁶ $T'_c \sim 10T'_*$, we expect that the total specific-heat of the double-layer, $C_v(T')$, has two smooth peaks in the regime $T'_* < T' < T'_c$.

N-layers. Let us now generalize the previous discussion to the case of N weakly coupled XY models (1). To simplify matters, let us also consider the partition function (2) in the absence of probe charges, i.e., $p = 0$. The dual in-plane field can then be expressed as $\vec{n}(\vec{r}, l) = \vec{n}'(\vec{r}, l) - \vec{n}_-(\vec{r}, l) + \vec{n}_-(\vec{r}, l-1)$, along with constraints $\vec{\nabla} \cdot \vec{n}' = 0$ and $\vec{\nabla} \cdot \vec{n}_- = n_z$. Again, taking the customary potential form for the “difference” field, $\vec{n}_- = -\vec{\nabla}\Phi$, and making suitable (lattice) integration by parts, one obtains the factorization $Z = Z_{\text{CG}} \Pi_{l=1}^{N-1} Z_{\text{DG}}^{(l)}$ for the partition function in the limit $J_{\perp}/J_{\parallel} \rightarrow 0$, where the Coulomb gas portion (4) now has the generalized form

$$\begin{aligned} Z_{\text{CG}} &= \sum_{\{n_z(\vec{r}, l)\}} \exp \left\{ -\frac{1}{2\beta_{\parallel}} \sum_{l=1}^N \sum_{\vec{r}} [\vec{n}_-(\vec{r}, l-1) - \vec{n}_-(\vec{r}, l)]^2 - \frac{1}{2\beta_{\perp}} \sum_{l=1}^{N-1} \sum_{\vec{r}} n_z^2(\vec{r}, l) \right\} \\ &= \sum_{\{n_z(\vec{r}, l)\}} \exp \left\{ -\frac{1}{2\beta_{\parallel}} \sum_{l=1}^N \sum_{\vec{r}, \vec{r}'} [n_z(\vec{r}, l-1) - n_z(\vec{r}, l)] G^{(2)}(\vec{r} - \vec{r}') [n_z(\vec{r}', l-1) - n_z(\vec{r}', l)] \right\} \end{aligned}$$

$$- \frac{1}{2\beta_{\perp}} \sum_{l=1}^{N-1} \sum_{\vec{r}} n_z^2(\vec{r}, l) \Big\}, \quad (8)$$

with the fields at the boundary layers set to $n_z(\vec{r}, 0) = 0 = n_z(\vec{r}, N)$. The key difference between the present Coulomb gas ensemble and the elementary one associated with the double-layer case (4) is that consecutive vertical fields, $n_z(\vec{r}, l-1)$ and $n_z(\vec{r}, l)$, of equal value make no contribution to the energy functional. However, in the limit of weakly coupled planes, Eq. (8) indicates that the concentration of dipolar (fluxon) charge excitations between consecutive planes is exponentially low. This observation suggests that each set of consecutive layers decouple precisely like the isolated double-layer at T_* , in agreement with the results obtained by Korshunov in his approximate treatment of the corresponding problem of infinitely many layers.¹² At temperatures below T_* , we expect that consecutive layers will be locked, with each layer in turn attaining quasi long-range order at temperatures below $T_c \sim T_*/10$. In this picture, therefore, the superfluid BKT transition involves the entire layered structure, making it 3D-like.¹³

The above picture may fail, however, in the regime of moderately coupled to strongly coupled planes, where the assumption that dipolar (fluxon) charge excitations be dilute is no longer valid. This is certainly true for the case of the isotropic problem with infinitely many layers (3D XY model), where the decoupling transition necessarily must be driven to infinite temperature due to the occurrence of a *single* 3D phase transition in the system. Hikami and Tsuneto have argued that vortex rings traversing many layers have a minimum mean radius given by $r \sim (J_{\parallel}/J_{\perp})^{1/2}$ in units of the lattice constants.¹⁴ Therefore, comparison of the latter to the total number of layers, N , yields a minimum anisotropy condition $J_{\parallel}/J_{\perp} \gtrsim N^2$ that insures the existence of the BKT superfluid transition characteristic of weakly coupled double-layers. This indicates, however, that the XY model with infinitely many layers may *not* be in the same universality class as that of the weakly coupled double-layer, contrary to claims in the literature.¹²

In conclusion, we find that the superfluid transition exhibited by a finite number (two or greater) of weakly coupled square-lattice XY model layers in the dual Villain form is 3D-like¹³ while remaining in the same universality class as that of the 2D XY model.¹² This result suggests that the latter transition is driven not by the unbinding of isolated vortex-antivortex pairs in a single layer, which are in fact (linearly) confined,^{14,15} but by the unbinding of *lines* of vortex-antivortex pairs passing perpendicularly through all of the

layers. We also find an inter-layer decoupling transition in the normal state at much higher temperature.¹² Last, in terms of the dual problems of lattice anyon superconductors and spin-liquids,⁶ these results indicate that the essential nature of the respective superconductor/normal and spin-liquid/paramagnetic transitions found in 2D is preserved in the present case of weakly coupled layered structures.

Discussions with M. Gabay, M. Maley, J. Engelbrecht and S. Trugman are gratefully acknowledged. This work was performed under the auspices of the U.S. Department of Energy, and was supported in part by the Associated Western Universities and National Science Foundation grant DMR-9322427.

References

1. *The Physical Properties of High-Temperature Superconductors*, vol. 2, edited by D.M. Ginsberg (World Scientific, Singapore, 1990).
2. P.W. Anderson, *Science* **235**, 1196 (1987).
3. R.B. Laughlin, *Phys. Rev. Lett.* **60**, 2677 (1988); R.B. Laughlin, *Science* **242**, 525 (1988).
4. J.P. Rodriguez and B. Douçot, *Phys. Rev. B* **45**, 971 (1992).
5. N. Read and S. Sachdev, *Phys. Rev. Lett.* **62**, 1694 (1989); J.P. Rodriguez, *Phys. Rev. B* **41**, 7326 (1990); A. Sokol and D. Pines, *Phys. Rev. Lett.* **71**, 2813 (1993).
6. J.P. Rodriguez, Los Alamos Report # LA-UR-94-3002 (1994).
7. S.T. Chui and J.D. Weeks, *Phys. Rev. B* **14**, 4978 (1976).
8. P. Minnhagen, *Rev. Mod. Phys.* **59**, 1001 (1987).
9. J.V. José, L.P. Kadanoff, S. Kirkpatrick and D.R. Nelson, *Phys. Rev. B* **16**, 1217 (1977); C. Itzykson and J. Drouffe, *Statistical field theory*, vol. 1, chap. 4 (Cambridge Univ. Press, Cambridge, 1991).
10. A. Polyakov, *Phys. Lett.* **72B**, 477 (1978); L. Susskind, *Phys. Rev. D* **20**, 2610 (1979).
11. X.G. Wen, F. Wilczek, and A. Zee, *Phys. Rev. B* **39**, 11413 (1989).
12. S.E. Korshunov, *Europhys. Lett.* **11**, 757 (1990).
13. S.T. Chui and M.R. Giri, *Phys. Lett. A* **128**, 49 (1988).
14. S. Hikami and T. Tsuneto, *Prog. Theor. Phys.* **63**, 387 (1980).
15. L.I. Glazmann and A.E. Koshelev, *Zh. Eksp. Teor. Fiz.* **97**, 1371 (1990) [*Sov. Phys. JETP* **70**, 774 (1990)]; V. Cataudella and P. Minnhagen, *Physica C* **166**, 442 (1990); P. Minnhagen and P. Olsson, *Phys. Rev. B* **44**, 4503 (1991).